

Remarks on Murre's conjecture on Chow groups*

by

KEJIAN XU AND ZE XU

Abstract.

For certain product varieties, Murre's conjecture on Chow groups is investigated. In particular, it is proved that Murre's conjecture (B) is true for two kinds of fourfolds. Precisely, if C is a curve and X is an elliptic modular threefold over k (an algebraically closed field of characteristic 0) or an abelian variety of dimension 3, then Murre's conjecture (B) is true for the fourfold $X \times C$.

Key Words: motivic decomposition, Chow group, curve, abelian variety, elliptic modular threefold

Mathematics Subject Classification 2010: 14C25

1. Introduction

We will work with the category \mathcal{V}_k of smooth projective varieties over a field k . Let $X \in \mathcal{V}_k$ be irreducible and of dimension d . Let $H(X) := H_{et}^*(\bar{X}, \mathbb{Q}_l)$ be the l -adic cohomology groups over a (fixed) algebraic closure \bar{k} of k , where $\bar{X} = X \times_k \text{Spec}(\bar{k})$ and $l \neq \text{ch}(k)$ is a prime, and let $\text{cl}_X : Z^i(X) \rightarrow H^{2i}(X)$ be the cycle map associated to $H(X)$, where $Z^i(X)$ is the group of algebraic cycles of codimension i of X . We have the well-known Künneth formula:

$$H^{2d}(X \times X) \simeq \bigoplus_{i=0}^{2d} H^{2d-i}(X) \otimes H^i(X).$$

Let $\Delta_X \subseteq X \times X$ be the diagonal. Then $\text{cl}_{X \times X}(\Delta_X)$ has the Künneth decomposition:

$$\text{cl}_{X \times X}(\Delta_X) = \pi_0^{\text{hom}} + \pi_1^{\text{hom}} + \dots + \pi_{2d}^{\text{hom}},$$

where $\pi_i^{\text{hom}} \in H^{2d-i}(X) \otimes H^i(X)$ is the i -th Künneth component.

Let $A_{\text{num}}^j(X)$ (resp. $A_{\text{rat}}^j(X) = \text{CH}^j(X)$) be the groups of algebraic cycles of codimension j modulo the numerical equivalence (resp. rational equivalence). Grothendieck's Lefschetz standard conjecture implies the π_i^{hom} are all algebraic (i.e., they are all in the image of the cycle map). Assuming additionally the conjecture that the homological equivalence coincides with the numerical equivalence ([13]), the diagonal (modulo the numerical equivalence) has a canonical decomposition into a sum of orthogonal idempotents (also called projectors)

$$\Delta_X = \pi_0^{\text{num}} + \pi_1^{\text{num}} + \dots + \pi_{2d}^{\text{num}}. \quad (1)$$

in the correspondence ring $A_{\text{num}}^d(X \times X) \otimes_{\mathbb{Z}} \mathbb{Q}$. Then, in the category of Grothendieck motives $\mathcal{M}_k^{\text{num}}$ ([13]) (w.r.t. the numerical equivalence), the motive $h(X) \in \mathcal{M}_k^{\text{num}}$ has a canonical decomposition

$$h(X) = h^0(X) \oplus h^1(X) \oplus \dots \oplus h^{2d}(X), \quad (2)$$

* This research is supported by National Natural Foundation of China (10871106).

where $h^i(X) := h(X, \pi_i^{\text{num}}, 0) \in \mathcal{M}_k^{\text{num}}$ (See [13] for details).

Furthermore, Murre ([15]) expected that the conjectural decomposition (1) exists even in $\text{CH}^d(X \times X; \mathbb{Q}) := \text{CH}^d(X \times X) \otimes_{\mathbb{Z}} \mathbb{Q} := A_{\text{rat}}^d(X \times X) \otimes_{\mathbb{Z}} \mathbb{Q}$ and hence in the category of Chow motives $\mathcal{M}_k^{\text{rat}}$ (w.r.t. the rational equivalence), $h(X) \in \mathcal{M}_k^{\text{rat}}$ has a decomposition as in (2). In this new setting, the decomposition is not canonical any more. However, from this conjectural decomposition, Murre ([15]) conjectured a very interesting filtration on rational Chow groups which relates the rational equivalence to the homological equivalence in finite steps as done by the conjectural Bloch-Beilinson filtration.

More precisely, as in [15], we will say that X has a *Chow-Künneth decomposition* over k if there exist $\pi_i \in \text{CH}^d(X \times X; \mathbb{Q})$, $0 \leq i \leq 2d$, satisfying

- (i) π_i are mutually orthogonal projectors;
- (ii) $\sum_i \pi_i = \text{cl}_{X \times X}(\Delta_X)$;
- (iii) $\text{cl}_{X \times X}(\pi_i) = \pi_i^{\text{hom}}$ (the i -th Künneth component).

Equivalently, the Chow motive of X has a (Chow-Künneth) decomposition

$$h(X) = h^0(X) \oplus h^1(X) \oplus \dots \oplus h^{2d}(X),$$

where $h^i(X) := h(X, \pi_i, 0) \in \mathcal{M}_k^{\text{rat}}$.

Then, Murre proposed in [15] the following famous conjecture.

Murre's Conjecture

(A): There exists a Chow-Künneth decomposition for every irreducible variety $X \in \mathcal{V}_k$ of dimension d .

(B): π_0, \dots, π_{j-1} and $\pi_{2j+1}, \dots, \pi_{2d}$ act as zero on $\text{CH}^j(X; \mathbb{Q})$.

(C): Let $F^v \text{CH}^j(X; \mathbb{Q}) = \text{Ker} \pi_{2j} \cap \text{Ker} \pi_{2j-1} \cap \dots \cap \text{Ker} \pi_{2j-v+1}$. Then the filtration F^\bullet is independent of the ambiguity in the choice of the π_i .

(D): $F^1 \text{CH}^j(X; \mathbb{Q}) = \text{CH}_{\text{hom}}^j(X; \mathbb{Q}) := \text{Ker}(\text{cl}_X)$.

It was shown by Jannsen ([9]) that Murre's conjecture is equivalent to the Bloch-Beilinson conjecture on rational Chow groups and the two conjectural filtrations proposed respectively by Murre, and Bloch and Beilinson coincides. The main advantage of Murre's conjecture over Bloch-Beilinson's is that one can check the statements for specific varieties as we will do in this paper.

Until now, Murre's conjecture is verified for only a few special varieties. It is known that (A) is true for curves, surfaces ([14]), Abelian varieties ([17][4]), Brauer-Severi varieties, some threefold ([2][3]), some special fourfold ([11]), certain modular varieties ([6][7]) and varieties whose Chow motives are finite-dimensional ([10]). As for the other parts of Murre's conjecture, it is known that (B) and (D) are true for the product of a curve and a surface ([15]), that (B) is true for the product of two surfaces ([11]) and that some part of (D) is true for the product of two surfaces ([11][12]). Jannsen ([10]) proved that (A), (B), (C) and (D) are true for some very special higher dimensional varieties over some special ground fields, in particular, he proved that if k is a rational or elliptic function field (in one variable) over a finite field \mathbb{F} and X_0 is an arbitrary product of rational and elliptic curves over \mathbb{F} , then (A)-(D) hold for $X_0 \times_{\mathbb{F}} k$. Gordon and Murre ([8]) proved that (A)-(D) are true for elliptic modular threefold over a field of characteristic 0.

In this paper, we consider Murre's conjecture for certain product varieties. Concretely, we consider such a problem: if the conjecture is true for X , when is it also true for the product of X with a curve or some other variety? In section 2, we consider the case of the product of a variety with a projective space. In section 3, we consider the case of the product of a variety with a curve. In particular, we generalize Murre's discussion given in [16], and as consequences, we prove that if C is a (smooth projective connected) curve, then Murre's conjecture (B) is true for $X \times C$, where X is an elliptic modular threefold over k (an algebraically closed field of characteristic 0) or X is an abelian variety of

dimension 3. This implies particularly that (B) is true for two new kinds of fourfolds other than products of two surfaces considered in [11] and [12].

2. Products with projective spaces

Fix a field k . Let X (resp. C) be a smooth projective irreducible variety (resp. curve) over k . Let X be of dimension d . In the following, we will always denote by $Z \in \mathrm{CH}^j(X)$ a cycle class. In addition, we denote by p with some lower indices the projection from a product variety to the corresponding factors.

In the proof of Theorem 2.3, the following lemma is crucial.

Lemma 2.1 ([5]) *Let \mathcal{E} be a vector bundle of rank $r = e + 1$ on a scheme X of finite type over $\mathrm{Spec}(k)$, with the projection $\pi : \mathcal{E} \rightarrow X$. Let $\mathbb{P}(\mathcal{E})$ be the associated projective bundle, p the projection from $\mathbb{P}(\mathcal{E})$ to X , and $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ the tautological line bundle on $\mathbb{P}(\mathcal{E})$. Then there are canonical isomorphisms*

$$\bigoplus_{i=0}^e \mathrm{CH}^{j-i}(X) \longrightarrow \mathrm{CH}^j(\mathbb{P}(\mathcal{E}))$$

$$(\alpha_i) \mapsto \sum_{i=0}^e c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))^i \cap p^* \alpha_i,$$

where $c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))$ is the first Chern class. □

Applying Proposition 3.1 in [5], it is easy to show that the inverse of the map in Lemma 2.1 is the map

$$\mathrm{CH}^j(\mathbb{P}(\mathcal{E})) \longrightarrow \bigoplus_{i=0}^e \mathrm{CH}^{j-i}(X), \quad \beta \mapsto (\beta_i),$$

where $\beta_e = p_* \beta$ and for $0 \leq i \leq e - 1$,

$$\beta_i = p_*(c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))^{e-i} \cap \beta) - \sum_{t=1}^{e-i} c_1(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))^{e+t} \cap p^* \beta_{i+t}.$$

Lemma 2.2 ([15]) *Assume that Y_i ($i = 1, 2$) are smooth projective irreducible varieties over k . Let $Y = Y_1 \times Y_2$. If Y_i ($i = 1, 2$) has a Chow-Künneth decomposition, then Y has also a Chow-Künneth decomposition.* □

Theorem 2.3 *Let X be a smooth projective irreducible variety of dimension d over k . If (A), (B) and (D) are true for X , then they are also true for $X \times \mathbb{P}^r$.*

Proof. Let $c = c_1(\mathcal{O}_{\mathbb{P}^r}(1))$, that is, the class of any hyperplane in \mathbb{P}^r . For each $0 \leq i \leq r$, set

$$\pi_{2i} = c^{r-i} \times c^i, \quad \pi_{2i+1} = 0, \quad h^i(\mathbb{P}^r) = (\mathbb{P}^r, \pi_i).$$

Then it is easy to see that there is the Chow-Künneth decomposition

$$h(\mathbb{P}^r) = \bigoplus_{i=0}^{2r} h^i(\mathbb{P}^r) = \bigoplus_{t=0}^r h^{2t}(\mathbb{P}^r).$$

Assume that X has the Chow-Künneth decomposition

$$h(X) = \bigoplus_{i=0}^{2d} h^i(X), \quad h^i(X) = (X, \pi'_i).$$

Then, $X \times \mathbb{P}^r$ has a Chow-Künneth decomposition:

$$h(X \times \mathbb{P}^r) = \bigoplus_{m=0}^{2(d+r)} h^m(X \times \mathbb{P}^r, \pi_m),$$

where $\pi_m = \sum_{p+2q=m} \tau_*(\pi'_p \times c^{r-q} \times c^q)$.

On the other hand, from Lemma 2.1, we have the isomorphisms:

$$\phi : \mathrm{CH}^j(X \times \mathbb{P}^r; \mathbb{Q}) \rightarrow \bigoplus_{i=0}^r \mathrm{CH}^{j-i}(X; \mathbb{Q}), \quad Z \mapsto (Z_i),$$

where $Z_i = p_{1*}([X] \times c^{r-i} \cdot Z)$ with

$$\phi^{-1}((Z_i)) = \sum_{i=0}^r ([X] \times c^i) \cdot p_1^* Z_i = \sum_{i=0}^r Z_i \times c^i,$$

and

$$\begin{aligned} \varphi : \mathrm{CH}^{d+r}(X \times \mathbb{P}^r \times X \times \mathbb{P}^r; \mathbb{Q}) &\longrightarrow \bigoplus_{i=0}^r \bigoplus_{t=0}^r \mathrm{CH}^{d+r-i-t}(X \times X; \mathbb{Q}), \\ \alpha &\longmapsto (\alpha_{it}) \end{aligned}$$

where $\alpha_{it} = p_{12*}([X \times X] \times c^{r-t} \times c^{r-i} \cdot \tau^* \alpha)$. In fact, we have

$$\begin{aligned} \alpha_{it} &= p_{13*}([X] \times c^{r-t} \times [X]) \cdot p_{123*}([X \times \mathbb{P}^r \times X] \times c^{r-i} \cdot \alpha) \\ &= p_{13*}([X] \times c^{r-t} \times [X] \times c^{r-i} \cdot \alpha) \\ &= p_{12*}([X \times X] \times c^{r-t} \times c^{r-i} \cdot \tau^* \alpha), \end{aligned}$$

where τ is the isomorphism exchanging the second and the third factor of the product variety $X \times X \times \mathbb{P}^r \times \mathbb{P}^r$. Clearly, we also have $\varphi^{-1}((\alpha_{it})) = \sum_{i,t} \tau_*(\alpha_{it} \times c^t \times c^i)$.

Now, define the map

$$\begin{aligned} \Phi : \mathrm{CH}^{d+r}(X \times \mathbb{P}^r \times X \times \mathbb{P}^r; \mathbb{Q}) \times \mathrm{CH}^j(X \times \mathbb{P}^r; \mathbb{Q}) &\longrightarrow \mathrm{CH}^j(X \times \mathbb{P}^r; \mathbb{Q}), \\ \Phi(\alpha, Z) &:= \alpha(Z) := p_{34*}(\alpha \cdot (Z \times [X \times \mathbb{P}^r])). \end{aligned}$$

So we have the following diagram

$$\begin{array}{ccc} \mathrm{CH}^{d+r}(X \times \mathbb{P}^r \times X \times \mathbb{P}^r; \mathbb{Q}) \times \mathrm{CH}^j(X \times \mathbb{P}^r; \mathbb{Q}) & \xrightarrow{\Phi} & \mathrm{CH}^j(X \times \mathbb{P}^r; \mathbb{Q}) \\ \downarrow \varphi \times \phi & & \downarrow \phi \\ \bigoplus_{i=0}^r \bigoplus_{t=0}^r \mathrm{CH}^{d+r-i-t}(X \times X; \mathbb{Q}) \times \bigoplus_{i=0}^r \mathrm{CH}^{j-i}(X; \mathbb{Q}) & \longrightarrow & \bigoplus_{i=0}^r \mathrm{CH}^{j-i}(X; \mathbb{Q}) \end{array}$$

Here the lower arrow is defined by the other three.

Note that if $t+i=r$, then we have

$$\begin{aligned} \tau_*(\pi'_p \times c^t \times c^q) \cdot (Z_i \times c^i) &= p_{34*}(\tau_*(\pi'_p \times c^t \times c^q) \cdot (Z_i \times c^i \times [X \times \mathbb{P}^r])) \\ &= p_{34*}\tau_*((\pi'_p \times c^t \times c^q) \cdot (Z_i \times [X] \times c^i \times [\mathbb{P}^r])) \\ &= p_{24*}((\pi'_p \cdot (Z_i \times [X])) \times c^{t+i} \times c^q) \\ &= p_{2*}((\pi'_p \cdot (Z_i \times [X])) \times c^{t+i}) \times c^q \\ &= \pi'_p(Z_i) \times c^q. \end{aligned}$$

So we conclude that

$$\tau_*(\pi'_p \times c^t \times c^q) \cdot (Z_i \times c^i) = \begin{cases} \pi'_p(Z_i) \times c^q, & \text{if } t+i=r; \\ 0, & \text{otherwise.} \end{cases}$$

Hence, we can translate the projectors on $X \times \mathbb{P}^r$ to those on X as follows.

$$\begin{aligned}
\phi(\pi_m(Z)) &= \phi \cdot \Phi \cdot (\varphi \times \phi)^{-1}(\varphi(\pi_m), (Z_i)) = \phi \cdot \Phi(\pi_m, \sum_{l=0}^r Z_l \times c^l) \\
&= \sum_{i=0}^r \sum_{p+2q=m} \phi(\tau_*(\pi'_p \times c^{r-q} \times c^q) \cdot (Z_i \times c^i)) \\
&= \sum_{i=0}^r \sum_{p+2q=m, i=q} \phi(\pi'_p(Z_i) \times c^q) \\
&= (\pi'_m(Z_0), \pi'_{m-2}(Z_i), \dots, \pi'_{m-2r}(Z_r))
\end{aligned}$$

Now, we can prove that the conjectures are true for $X \times \mathbb{P}^r$.

For (B), from $0 \leq m \leq j-1$ we have $m-2i \leq (j-i)-1$. If $2j+1 \leq m \leq 2(d+r)$, then

$$2(j-i)+1 \leq m-2i \iff 2j+1 \leq m.$$

So, by the assumptions on X , we see that

$$\pi_m(Z) = 0, \quad \text{for } 0 \leq m \leq j-1 \text{ or } 2j+1 \leq m.$$

For (D), suppose that $Z \in \text{CH}_{\text{hom}}^j(X \times \mathbb{P}^r; \mathbb{Q}) = \text{Ker}(\text{cl}_{X \times \mathbb{P}^r})$. Then since $Z_i \in \text{Ker}(\text{cl}_X) = \text{CH}_{\text{hom}}^{j-i}(X; \mathbb{Q}) = \text{Ker}(\pi'_{2(j-i)})$ by assumption, we have $Z \in \text{Ker}(\pi_{2j})$. This completes the proof of Theorem 2.3. \square

Remark 2.4 (i) We expect that Theorem 2.3 is also true for non-trivial projective bundles.

(ii) For (C), we can say nothing yet since from the projectors on X we can get only one but not all set of projectors on $X \times \mathbb{P}^r$.

Corollary 2.5 *Let S_1, S_2 be smooth projective surfaces over k . Then conjectures (A) and (B) are true for $S_1 \times S_2 \times \mathbb{P}^{r_1} \times \dots \times \mathbb{P}^{r_n}$.*

Proof: From Lemma 2.2 and the main theorem of [14], we know that conjectures (A) and (B) are true for $S_1 \times S_2$, so the result follows from Theorem 2.3. \square

3. Products with curves

Let C be a smooth projective curve over a field k and $e \in C(k)$. It is well-known (see [18] for details) that C has the Chow-Künneth decomposition

$$h(C) = h^0(C) \oplus h^1(C) \oplus h^2(C),$$

where $h^i(C) = (C, \pi_i'')$ with

$$\pi_0'' = [e \times C], \quad \pi_2'' = [C \times e], \quad \pi_1'' = \Delta_C - \pi_0'' - \pi_2''.$$

Assume that the irreducible variety $X \in \mathcal{V}_k$ has the Chow-Künneth decomposition

$$h(X) = \bigoplus_{i=0}^{2d} h^i(X), \quad h^i(X) = (X, \pi_i').$$

Then, the product variety $X \times C$ has the Chow-Künneth decomposition

$$h(X \times C) = \bigoplus_{m=0}^{2(d+1)} h^m(X \times C, \pi_m),$$

where, explicitly,

$$\pi_0 = \pi_0' \times [e \times C],$$

$$\begin{aligned}\pi_1 &= \pi'_1 \times [e \times C] + \pi'_0 \times (\Delta_C - [e \times C] - [C \times e]), \\ \pi_m &= \pi'_m \times [e \times C] + \pi'_{m-1} \times (\Delta_C - [e \times C] - [C \times e]) + \pi'_{m-2} \times [C \times e], \quad m \geq 2.\end{aligned}$$

Let

$$\mathrm{CH}_{\mathrm{alg}}^j(X; \mathbb{Q}) := \{Z \in \mathrm{CH}^j(X; \mathbb{Q}) : Z \sim_{\mathrm{alg}} 0\},$$

where $Z \sim_{\mathrm{alg}} 0$ means that Z is algebraically equivalent to 0.

In the proof of Theorem 3.3, we need the following computations.

Lemma 3.1 *For any $Z \in \mathrm{CH}^j(X \times C; \mathbb{Q})$, we have*

- (i) $(\pi'_m \times [e \times C])(Z) = \pi'_m(Z(e)) \times [C]$;
- (ii) $(\pi'_m \times [C \times e])(Z) = \pi'_m(p_{1*}Z) \times [e]$.

Proof: (i) We have

$$\begin{aligned}(\pi'_m \times [e \times C])(Z) &= p_{34*}(\tau_*(\pi'_m \times [e \times C]) \cdot (Z \times [X \times C])) \\ &= p_{34*}(\tau_*(\pi'_m \times [e]) \cdot (Z \times [X]) \times [C]) \\ &= p_{3*}(\tau_*(\pi'_m \times [e]) \cdot (Z \times [X]) \times [C]) \\ &= \pi'_m(Z(e)) \times [C],\end{aligned}$$

where $Z(e) = p_{1*}(Z \cdot (X \times [e]))$. Note that in the last equality, we have used the following computation.

$$\begin{aligned}p_{3*}(\tau_*(\pi'_m \times [e]) \cdot (Z \times [X])) &= p_{3*}(\tau_*((\pi'_m \times [C]) \cdot [X \times [e] \times X) \cdot (Z \times [X])) \\ &= p_{3*}((p_{13}^* \pi'_m \cdot [X \times e \times X]) \cdot (Z \times [X])) \\ &= p_{3*}((p_{13}^* \pi'_m \cdot ((Z \cdot [X \times e]) \times [X])) \\ &= p_{2*}p_{13*}((p_{13}^* \pi'_m \cdot ((Z \cdot [X \times e]) \times [X])) \\ &= p_{2*}(\pi'_m \cdot p_{13*}((Z \cdot [X \times e]) \times [X])) \\ &= p_{2*}(\pi'_m \cdot (p_{13*}(Z \cdot [X \times e]) \times [X])) \\ &= \pi'_m(Z(e)).\end{aligned}$$

(ii) Similar to (i), we have

$$\begin{aligned}(\pi'_m \times [C \times e])(Z) &= p_{34*}(\tau_*(\pi'_m \times [C \times e]) \cdot (Z \times [X \times C])) \\ &= p_{34*}(\tau_*(\pi'_m \times [C]) \cdot (Z \times [X]) \times [e]) \\ &= p_{3*}((\tau_*(\pi'_m \times [C]) \cdot (Z \times [X])) \times [e]) \\ &= \pi'_m(p_{1*}Z) \times [e].\end{aligned}$$

□

Lemma 3.2 *Let $Z \in \mathrm{CH}^j(X \times C; \mathbb{Q})$. Then*

- (i) $(\pi'_m \times \Delta_C)(Z) = p_{23*}(p_{13}^*Z \cdot (\pi'_m \times [C]))$;
- (ii) $(\mathrm{id}_X \times f)^*((\pi'_m \times \Delta_C)(Z)) = (\pi'_m)_K((\mathrm{id}_X \times f)^*Z)$, where $K = k(C)$ is the function field of C , $f : \mathrm{Spec}(K) \rightarrow C$ is the natural morphism and $(\pi'_m)_K = \pi'_m \times \Delta_{\mathrm{Spec}(K)}$.

Proof: (i) Let $\delta_C : C \rightarrow C \times C$ be diagonal morphism. Then, we have

$$\begin{aligned}(\pi'_m \times \Delta_C)(Z) &= p_{34*}(\tau_*(\pi'_m \times \Delta_C) \cdot (Z \times [X \times C])) \\ &= p_{34*}(\tau_*(\mathrm{id}_{X \times X} \times \delta_C)_*(\pi'_m \times \Delta_C) \cdot (Z \times [X \times C])) \\ &= p_{23*}((\pi'_m \times \Delta_C) \cdot (\mathrm{id}_{X \times X} \times \delta_C)^* \tau^*Z) \\ &= p_{23*}(p_{13}^*Z \cdot (\pi'_m \times [C])).\end{aligned}$$

(ii) From (i) and the following diagram

$$\begin{array}{ccc} X_K \times_K X_K & \xrightarrow{p_{X_K}} & X_K \\ \mathrm{id}_X \times \mathrm{id}_X \times f \downarrow & & \downarrow \mathrm{id}_X \times f \\ X \times X \times C & \xrightarrow{p_{23}} & X \times C \end{array}$$

we have

$$\begin{aligned}
(\mathrm{id}_X \times f)^*((\pi'_m \times \Delta_C)(Z)) &= (\mathrm{id}_X \times f)^* p_{23*}(p_{13}^* Z \cdot \pi'_m \times [C]) \\
&= p_{X_K*}(\mathrm{id}_X \times \mathrm{id}_X \times f)^*(p_{13}^* Z \cdot \pi'_m \times [C]) \\
&= p_{X_K*}((\mathrm{id}_X \times \mathrm{id}_X \times f)^* p_{13}^* Z \cdot (\mathrm{id}_X \times \mathrm{id}_X \times f)^*(\pi'_m \times [C])) \\
&= p_{X_K*}(((\mathrm{id}_X \times f)^* Z \times_K X_K) \cdot \pi'_m \times \Delta_{\mathrm{spec} K}) \\
&= (\pi'_m \times \Delta_{\mathrm{spec} K})((\mathrm{id}_X \times f)^* Z) \\
&= (\pi'_m)_K((\mathrm{id}_X \times f)^* Z).
\end{aligned}$$

□

Our main theorem is the following

Theorem 3.3 *Let k be an algebraically closed field, $X \in \mathcal{V}(k)$ and $C \in \mathcal{V}(k)$ an irreducible curve with the function field $K = k(C)$. Assume that (A) and (B) are true for X and X_K , and that for any j , $\mathrm{CH}_{\mathrm{alg}}^j(X_K; \mathbb{Q}) \subseteq \mathrm{Ker}((\pi'_{2j})_K)$. Then (A) and (B) are also true for $X \times C$.*

Proof: The statement about (A) is obvious. We will consider (B) in the following. Let $Z \in \mathrm{CH}^j(X \times C; \mathbb{Q})$. Easy computations shows that (B) is true if Z is of the form $Z' \times [C]$ with $Z' \in \mathrm{CH}^j(X; \mathbb{Q})$. So, we can assume $Z(e) = 0$, since we have

$$\begin{aligned}
Z &= (Z - Z(e) \times [C]) + Z(e) \times [C], \\
(Z - Z(e) \times [C])(e) &= Z(e) - Z(e) = 0.
\end{aligned}$$

Assume that

$$0 \leq m \leq j-1 \quad \text{or} \quad 2j+1 \leq m \leq 2(d+1).$$

Then, for $m \geq 1$, we have

$$0 \leq m-1 \leq (j-1)-1 \quad \text{or} \quad 2(j-1)+1 \leq m-1,$$

and for $m \geq 2$, we have

$$0 \leq m-2 \leq (j-1)-1 \quad \text{or} \quad 2(j-1)+1 \leq m-1.$$

From Lemma 3.1 and the assumptions on X , we have (note that $p_{1*}Z \in \mathrm{CH}^{j-1}(X; \mathbb{Q})$)

$$\begin{aligned}
(\pi'_m \times [e \times C])(Z) &= \pi'_m(Z(e)) \times [C] = 0, \\
(\pi'_{m-1} \times [e \times C])(Z) &= \pi'_{m-1}(Z(e)) \times [C] = 0, \\
(\pi'_{m-1} \times [C \times e])(Z) &= \pi'_{m-1}(p_{1*}Z) \times [e] = 0, \\
(\pi'_{m-2} \times [C \times e])(Z) &= \pi'_{m-2}(p_{1*}Z) \times [e] = 0.
\end{aligned}$$

So, the problem is reduced to prove

$$(\pi'_{m-1} \times \Delta_C)(Z) = 0, \quad \text{if } 1 \leq m \leq j-1 \text{ or } 2j+1 \leq m \leq 2(d+1).$$

At first, we show that $(\mathrm{id} \times f)^*Z$ is algebraically equivalent to 0 on X_K . In fact, let η be the generic point of C , that is, $K = k(\eta)$, and let $f_\eta : \mathrm{Spec}(K) \rightarrow C_K$ be the K -point defined by η . Denote $\eta_K = f_\eta(\mathrm{Spec} K)$. Then we have

$$\begin{aligned}
Z_K(\eta_K) &= p_{X_K*}(Z_K \cdot X \times \eta_K) = p_{X_K*}(Z_K \cdot (\mathrm{id}_X \times f_\eta)_*(X \times \mathrm{Spec} K)) \\
&= p_{X_K*}(\mathrm{id}_X \times f_\eta)_*((\mathrm{id}_X \times f_\eta)^* Z_K) = (\mathrm{id}_X \times f_\eta)^* Z_K \\
&= (\mathrm{id}_X \times f_\eta)^* p_{X_K}^{XCK*}(Z) = (\mathrm{id}_X \times f)^*(Z).
\end{aligned}$$

Similarly, let $g_e : \mathrm{Spec}(K) \rightarrow C_K$ and $g : \mathrm{Spec} k \rightarrow C$ be the morphisms both defined by e . Denote $e_K = g_e(\mathrm{Spec}(K))$. Then we have $Z_K(e_K) = ((\mathrm{id}_X \times g)^*(Z))_K = Z(e)_K = 0$.

We claim that

$$(\pi'_{m-1})_K((\mathrm{id} \times f)^*Z) = 0, \quad \text{if } 1 \leq m \leq j-1 \text{ or } 2j+1 \leq m \leq 2(d+1).$$

In fact, if $m = 2j + 1$, since $(\text{id} \times f)^*Z \in \text{CH}^j(X_K; \mathbb{Q})$ is algebraically equivalent to 0, from the assumption $\text{CH}_{\text{alg}}^j(X_K; \mathbb{Q}) \subseteq \text{Ker}((\pi'_{2j})_K)$ we have

$$(\pi'_{m-1})_K((\text{id} \times f)^*Z) = (\pi'_{2j})_K((\text{id} \times f)^*Z) = 0;$$

if $1 \leq m \leq j - 1$ or $2j + 2 \leq m \leq 2(d + 1)$, we have $1 \leq m - 1 \leq j - 2$ or $2j + 1 \leq m - 1 \leq 2d + 1$, so from the assumptions on X we get $(\pi'_{m-1})_K((\text{id} \times f)^*Z) = 0$ since $(\text{id} \times f)^*Z \in \text{CH}^j(X_K; \mathbb{Q})$.

On the other hand, we have the following well known diagram

$$\begin{array}{ccccccc} \text{CH}^{j-1}(X \times (C - U); \mathbb{Q}) & \longrightarrow & \text{CH}^j(X \times C; \mathbb{Q}) & \longrightarrow & \text{CH}^j(X \times U; \mathbb{Q}) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ & & \text{CH}^j(X_K; \mathbb{Q}) & \xlongequal{\quad} & \lim_{U \subseteq C} \text{CH}^j(X \times U; \mathbb{Q}) & & \end{array}$$

where the left vertical map is $z \mapsto (\text{id} \times f)^*z$. So from Lemma 3.2 (ii), we have

$$(\text{id}_X \times f)^*((\pi'_{m-1} \times \Delta_C)(Z)) = (\pi'_{m-1})_K((\text{id} \times f)^*Z) = 0,$$

hence

$$(\pi'_{m-1} \times \Delta_C)(Z) = \sum_i Z'_i \times a_i, \text{ with } Z'_i \in \text{CH}^{j-1}(X; \mathbb{Q}) \text{ and } a_i \in \text{CH}^1(C; \mathbb{Q}).$$

In view of $(\pi'_{m-1} \times \Delta_C)^2 = \pi'_{m-1} \times \Delta_C$, we conclude that

$$\begin{aligned} (\pi'_{m-1} \times \Delta_C)(Z) &= \sum_i (\pi'_{m-1} \times \Delta_C)(Z'_i \times a_i) \\ &= \sum_i p_{23*}[p_{13}^*(Z'_i \times a_i) \cdot (\pi'_{m-1} \times [C])] \\ &= \sum_i \pi'_{m-1}(Z'_i) \times a_i = 0. \end{aligned}$$

This completes the proof of the theorem. \square

Although the theorem above is restricted, we can deduce several interesting consequences.

Corollary 3.4 *If k is an algebraically closed field of characteristic 0 and X is an elliptic modular threefold over k , then (A) and (B) are true for $X \times C$.*

Proof: It was shown in [8] that Murre's conjecture holds for an elliptic modular threefold over a field of characteristic 0. Obviously, conjecture (D) for X_K implies the assumption of Theorem 3.3. So, the result is an immediate consequence of Theorem 3.3. \square

Corollary 3.5 *Assume that algebraic equivalence and rational equivalence coincide on X . If (A) and (B) are true for X and X_K , then (A) and (B) are also true for $X \times C$.* \square

Remark 3.6 Cellular varieties satisfy the first hypothesis of the corollary.

By [1] (see also [4] and [15]), for an abelian variety X of dimension g over any field k , we have the following decomposition

$$\text{CH}^j(X; \mathbb{Q}) = \bigoplus_{s=j-g}^j \text{CH}_s^j(X),$$

where

$$\text{CH}_s^j(X) := \{\alpha \in \text{CH}^j(X; \mathbb{Q}) \mid n^* \alpha = n^{2j-s} \alpha, \forall n \in \mathbb{Z}\}.$$

Corollary 3.7 *Let X be an abelian variety of dimension at most 4 over an algebraically closed field k . Assume that for any j , $CH_0^j(X_K) \cap CH_{\text{alg}}^j(X_K; \mathbb{Q}) = 0$. Then (A) and (B) are true for $X \times C$.*

Proof: It follows from [1] that conjecture (B) is true for an abelian variety of dimension at most 4, equivalently, Beauville's vanishing conjecture holds: $CH_s^j(X) = 0$ if $s < 0$. By assumption and the fact that the algebraic equivalence is adequate, we have

$$CH_{\text{alg}}^j(X_K; \mathbb{Q}) = \bigoplus_{s=1}^j (CH_s^j(X_K) \cap CH_{\text{alg}}^j(X_K; \mathbb{Q})).$$

On the other hand, by Lemma 2.5.1 of [15], we see that for any j ,

$$\text{Ker}((\pi'_{2j})_K) = \bigoplus_{s=1}^j CH_s^j(X_K).$$

Then we can apply Theorem 3.3 to end the proof. \square

Remark 3.8 The assumption of Corollary 3.7 is a consequence of a conjecture of Beauville: the restricted cycle map $c_0 : CH_0^j(X) \rightarrow H^{2j}(X)$ is injective for any j .

Corollary 3.9 *Let X be an abelian variety of dimension 3 over an algebraically closed field k . Then (A) and (B) are true for $X \times C$.*

Proof: By [1], we know that the restricted cycle map

$$c_0 : CH_0^j(X_K) \rightarrow H^{2j}(X_K)$$

is injective for $j = 0, 1, g-1, g$ if X is an abelian variety of dimension g . So for any j ,

$$CH_0^j(X_K) \cap CH_{\text{alg}}^j(X_K; \mathbb{Q}) \subseteq CH_0^j(X_K) \cap CH_{\text{hom}}^j(X_K; \mathbb{Q}) = 0.$$

Hence the result follows from Corollary 3.7. \square

REFERENCES

1. A. Beauville. Sur l'anneau de Chow d'une variété abélienne. *Math. Ann.* **273**(1986), 647-651
2. P. L. del Angel and S. Müller-Stach. Motives of uniruled 3-folds, *Compositio Math.* **112**(1998), 1-16
3. P. L. del Angel, S. Müller-Stach. On Chow motives of 3-folds. *Trans. Amer. Math. Soc.* **352**(2000), 1623-1633
4. C. Deninger and J. P. Murre. Motivic decomposition of abelian schemes and the Fourier transform. *J. Reine Angew. Math.* **422**(1991), 201-219
5. W. Fulton. Intersection theory. *Ergeb. Math. Grenzgeb.* Springer-Verlag, Berlin 1984
6. B. Gordon, M. Hanamura and J. P. Murre. Relative Chow-Künneth projectors for modular varieties. *J. reine angew. Math.* **558** (2003), 1-14
7. B. Gordon, M. Hanamura and J. P. Murre. Absolute Chow-Künneth projectors for modular varieties. *J. reine angew. Math.* **580**(2005), 139-155
8. B. Gordon and J. P. Murre. Chow motive of elliptic modular surfaces and threefolds. *Math. Inst., Univ. of Leiden*, Report W 96-16 (1996)
9. U. Jannsen. Motivic sheaves and filtrations on Chow groups. In *Proc. sympos. Pure Math.* **55**(1994), Part 1, 245-302
10. U. Jannsen. On finite-dimensional motive and Murre's conjecture. Preprint, 2007
11. B. Kahn, J. P. Murre and C. Pedrini. On the transcendental part of the motive of

- a surface. Preprint, 2005
12. K. Kimura. Murre's conjectures for certain product varieties. *J. Math. Kyoto. Univ.* **47**(3)(2007), 621-629
 13. J. P. Murre. Lectures on motives. In *Transcendental Aspects of Algebraic Cycles*, Proceedings of Grenoble Summer School(2001), pages 123-170, Cambridge University Press, 2004
 14. J. P. Murre. On the motive of an algebraic surface. *J. Reine und angew. Math.* **409**(1990), 190-204
 15. J. P. Murre. On a conjectural filtration on the Chow groups of an algebraic variety, I. The general conjectures and some examples. *Indag. Math. N. S.* **4**(2)(1993), 189-201
 16. J. P. Murre. On a conjectural filtration on the Chow groups of an algebraic variety, II. Verification of the conjectures for threefolds which are the product on a surface and a curve. *Indag. Math. N. S.* **4**(2)(1993), 189-201
 17. A. M. Shermenev. The motive of an abelian variety. *Funct. Anal.* **8**(1974), 47-53
 18. A.J. Scholl. Classical motives, In *Proceedings of Symposia in Pure Mathematics*, **55**(1994), Part I, 163-187

KEJIAN XU kejianxu@amss.ac.cn

College of Mathematics
Qingdao University
Qingdao 266071, China

ZE XU xuze@amss.ac.cn

Institute of Mathematics
Academy of Mathematics and System Science
Chinese Academy of Sciences
Beijing 100190, China